



Skip this form & e-prescribe! Select BioPlus from your EHR!

Fax: 866-523-5406
Phone: 800-829-3975
bioplusinfusion.com

IVIG

Ship To: [ ] In Office [ ] Infusion Suite [ ] At Home [ ] Other

PATIENT INFORMATION

Name: SSN: DOB:
Address: City: State: ZIP:
Home Phone: Cell: Height: Weight: Gender: Female Male
Email: Allergies:

INSURANCE INFORMATION (or attach copy of cards)

Primary Insurance Co: Policy Holder: Relationship: Policy #: Group #:
Secondary Insurance: Policy Holder: Relationship: Policy #: Group #:

CLINICAL INFORMATION (Fax all pertinent clinical and lab information)

Diagnosis (ICD-10): Date of Diagnosis:
Pemphigus L10.0 CIDP G61.81 Peripheral Neuropathy G60.9 MMN G61.82 Multiple Sclerosis G35 Acute Infective Polyneuritis/GBS G61.0
Myasthenia Gravis with acute exacerbation G70.01 Myasthenia Gravis without acute exacerbation G70.00 Dermatomyositis M33.90 Polymyositis M33.20
Has patient received immune globulin previously? [ ] No [ ] Yes: Date of last infusion Date of next infusion:
Comorbidities:
ALLERGIES: [ ] NKDA [ ] Other

PRESCRIPTION INFORMATION (or attach a copy of the prescription)

Infusion Therapy:

Preferred brand OR [ ] Pharmacist will determine appropriate product based on clinical assessment, insurance requirements, and availability
[ ] No Substitute [ ] Refills: times (as allowed by state or payer requirements)

Dose: (please select option(s) and provide complete information, pharmacy to round the nearest 5 gram vial)

- [ ] Administration Rate = Follow Manufacturer's Guidelines
[ ] Loading Dose: gm/kg over days, then
[ ] Maintenance dose: gm/kg over days, every weeks x cycles
[ ] Other Regimen

Infusion Rate: (please select one and provide complete information)

- [ ] Pharmacist to determine
[ ] Start at mL/hr, then increase by mL/hr every minutes to maximum rate mL/hr

Vascular Access Device:

- [ ] Peripheral Catheter [ ] PICC [ ] Port
[ ] Other (describe # of lumens):

Flush Orders: (If IV ordered, the following flush protocols will be followed)

- [ ] Sodium Chloride 0.9%
Peripheral Line: 3 mL before each dose and 3 mL after each dose and PRN
Central Line: 5-10 mL before each dose and 5-10 mL after each dose and PRN
[ ] Heparin 10 u/mL Peripheral Line: 3 mL after last sodium flush and PRN
[ ] Heparin 100 u/mL Central Line: 5 mL after last sodium flush and PRN
Provide needles, syringes, VAD supplies & other ancillary supplies needed for infusion

Hydration Orders

Infuse mg solution
[ ] Prior to [ ] Following

Labs:

Results will be faxed to physician's office. If no frequency noted, ordered labs to be done prior to initial infusion only. Labs will not be drawn on weekend/holidays. Not appropriate for STAT labs
[ ] Quantitative Ig A prior to first dispense. Pharmacist to obtain authorization from MD.
[ ] Other Frequency of Labs:

Anaphylaxis Kit Order (Infusion Reaction Management x 1/year)

[ ] Epinephrine [ ] IM [ ] SUBQ Qty: Refills:
Adult 1:1000, 0.3 mL (>30 kg/>66lbs) PRN severe allergic reaction, call 911. May repeat in 5-15 minutes, as needed.

As required by your state, Prescriber to check "Dispense as written" or handwritten "Brand Medically Necessary" and sign to prevent generic substitution. [ ] Dispense as written

NURSING

[ ] Nursing Agency: Phone:
Skilled Nursing Visits for Immune Globulin Intravenous administration and education. To provide education related to the disease process and IG therapy.
To provide an assessment of patient's general overall health status. To provide skilled nursing visits PRN for additional education and support.

PHYSICIAN INFORMATION

Prescriber Name: Phone: Fax:
Office Contact: Email:
Address: City: State: ZIP:
NPI #: Tax ID:
Prescriber Signature: Date: