

BLEEDING DISORDER

Ship To: In Office Infusion Suite At Home Other _____

PATIENT INFORMATION

Name:		SSN:	DOB:	
Address:		City:	State:	ZIP:
Home Phone:	Cell:	Height:	Weight:	Gender: Female Male
Email:		Allergies:		

INSURANCE INFORMATION (or attach copy of cards)

Primary Insurance:	Policy Holder:	Relationship:	Policy #:	Group #:
Secondary Insurance:	Policy Holder:	Relationship:	Policy#:	Group#:

CLINICAL INFORMATION

Diagnosis (ICD-10): _____

D66 Hereditary Factor VIII Disorder (Hemophilia A) Severity mild moderate severe
 D67 Hereditary Factor IX Disorder (Hemophilia B) Severity mild moderate severe
 D68.0 Von Willebrand Disease Type: 1 2A 2B 2M 2n 3
 D68.31 1 Acquired Hemophilia
 D68.9 Coagulation defect, unspecified

D68.2 Hereditary deficiency of other clotting factors
 D68.318 Other hemorrhagic disorder due to intrinsic circulating anticoagulants, antibodies, or inhibitors
 Other _____

Date of Diagnosis: _____ Comorbidities: _____

ALLERGIES: NKDA Other _____

PRESCRIPTION INFORMATION (or attach a copy of prescription)

CLOTTING FACTOR ORDERS

Brand Name: _____ Dose: _____ Qty: _____ Frequency: _____
 Brand Name: _____ Dose: _____ Qty: _____ Frequency: _____
 Dosage: Mild units/kg _____ Severe units/kg _____
 Prophylaxis: Dispense _____ dose/week for a duration of _____ months
 Episodic: Dispense _____ doses for mild or _____ doses for severe

OTHER MEDICATION

Amicar® Directions: _____ Qty: _____ Refills: _____
 Lysteda® Directions: _____ Qty: _____ Refills: _____
 Stimate® Directions: _____ Qty: _____ Refills: _____
 _____ Directions: _____ Qty: _____ Refills: _____

VASCULAR ACCESS DEVICE: Peripheral Catheter PICC Port Other (describe/# of lumens): _____

Flush Orders: (If IV ordered, the following flush protocols will be followed)

Sodium Chloride 0.9% Other _____

Peripheral Line: 3 mL before each dose and 3 mL after each dose and PRN

Central Line: 5-10 mL before each dose and 5-10 mL after each dose and PRN

Anaphylaxis Kit Order (Infusion Reaction Management x 1/year)

Epinephrine IM SUBQ Qty: _____ Refills: _____
 Adult 1:1000, 0.3 mL (>30 kg/>66lbs) PRN severe allergic reaction, call 911. May repeat in 5-15 minutes as needed.

NURSING

Nursing Agency: _____ Phone: _____
 Skilled Nursing Visits for infliximab intravenous administration and education. To provide education related to the disease process and therapy. To provide an assessment of patient's general overall health status. To provide skilled nursing visits PRN for additional education and support.

As required by your state, Prescriber to check "Dispense as written" or handwritten "Brand Medically Necessary" and sign to prevent generic substitution. Dispense as written

PHYSICIAN INFORMATION

INFUSION TYPE: At Home In Office

Prescriber Name:	Phone:	Fax:	
Office Contact:	Email:		
Address:	City:	State:	ZIP:
NPI #:	Tax ID:		
Prescriber Signature:	Date:		